

Amino acids and central fatigue.

[Blomstrand E.](#)

Department of Sport and Health Sciences, University College of Physical Education and Sports, Karolinska Institute, Stockholm, Sweden. eva.blomstrand@ihs.se

There is an increasing interest in the mechanisms behind central fatigue, particularly in relation to changes in brain monoamine metabolism and the influence of specific amino acids on fatigue. Several studies in experimental animals have shown that physical exercise increases the synthesis and metabolism of brain 5-hydroxytryptamine (5-HT). Support for the involvement of 5-HT in fatigue can be found in studies where the brain concentration of 5-HT has been altered by means of pharmacological agents. When the 5-HT level was elevated in this way the performance was impaired in both rats and human subjects, and in accordance with this a decrease in the 5-HT level caused an improvement in running performance in rats. The precursor of 5-HT is the amino acid tryptophan and the synthesis of 5-HT in the brain is thought to be regulated by the blood supply of free tryptophan in relation to other large neutral amino acids (including the branched-chain amino acids, BCAA) since these compete with tryptophan for transport into the brain. Studies in human subjects have shown that the plasma ratio of free tryptophan/BCAA increases during and, particularly, after sustained exercise. This would favour the transport of tryptophan into the brain and also the synthesis and release of 5-HT which may lead to central fatigue. Attempts have been made to influence the 5-HT level by giving BCAA to human subjects during different types of sustained heavy exercise. The results indicate that ingestion of BCAA reduces the perceived exertion and mental fatigue during exercise and improves cognitive performance after the exercise. In addition, in some situations ingestion of BCAA might also improve physical performance; during exercise in the heat or in a competitive race when the central component of fatigue is assumed to be more pronounced than in a laboratory experiment. However, more experiments are needed to further clarify the effect of BCAA and also of tryptophan ingestion on physical performance and mental fatigue.

Related articles

- [A role for branched-chain amino acids in reducing central fatigue.](#)

J Nutr. 2006 Feb; 136(2):544S-547S.

[J Nutr. 2006]

- [ReviewTryptophan, 5-hydroxytryptamine and a possible explanation for central fatigue.](#)

Adv Exp Med Biol. 1995; 384:315-20.

[Adv Exp Med Biol. 1995]

- [Changes in the albumin binding of tryptophan during postoperative recovery: a possible link with central fatigue?](#)

Brain Res Bull. 1997; 43(1):43-6.

[Brain Res Bull. 1997]

- [ReviewThe role of tryptophan in fatigue in different conditions of stress.](#)

Adv Exp Med Biol. 1999; 467:697-704.

[Adv Exp Med Biol. 1999]

- [Administration of branched-chain amino acids during sustained exercise--effects on performance and on plasma concentration of some amino acids.](#)

Eur J Appl Physiol Occup Physiol. 1991; 63(2):83-8.

[*Eur J Appl Physiol Occup Physiol. 1991*]

Cited by 3 PubMed Central articles

- [Dietary supplements and sports performance: amino acids.](#)

Williams M. J Int Soc Sports Nutr. 2005 Dec 9; 2:63-7. Epub 2005 Dec 9.

[*J Int Soc Sports Nutr. 2005*]

- [ReviewHow to increase serotonin in the human brain without drugs.](#)

Young SN. J Psychiatry Neurosci. 2007 Nov; 32(6):394-9.

[*J Psychiatry Neurosci. 2007*]

- [ReviewEvidence for complex system integration and dynamic neural regulation of skeletal muscle recruitment during exercise in humans.](#)

St Clair Gibson A, Noakes TD. Br J Sports Med. 2004 Dec; 38(6):797-806.

[*Br J Sports Med. 2004*]